

# *Ctenopteryx sicula*, a bathypelagic loliginid squid?

Andrew S. Brierley\*, Malcolm R. Clarke, and John P. Thorpe

Port Erin Marine Laboratory, Department of Environmental and Evolutionary Biology, University of Liverpool, Isle of Man, IM9 6JA, British Isles

**Abstract:** *Ctenopteryx sicula* (Vérany, 1851) is an open-eyed, deep-water squid, and as such has traditionally been classified in the suborder Oegopsida (family Ctenopterygidae) along with other families of squid exhibiting these characteristics. *C. sicula* however displays numerous morphological features, including fused axons in the giant nerve fiber system and accessory nidamental glands, found otherwise only within members of the myopsid families Loliginidae and Pickfordioteuthidae. This has led previous authors to suggest that *Ctenopteryx* species would be more appropriately placed in the suborder Myopsida. Here biochemical genetic evidence is presented which indicates that *C. sicula* is more closely related to several loliginid species than to species of the oegopsid families Histiototeuthidae, Ommastrephidae, and Enoploteuthidae. These data, in conjunction with new data on comparative beak morphology, also suggest that *C. sicula* should be considered an oceanic myopsid species.

The squids (Teuthoidea) are generally considered to be divided into two suborders: the Oegopsida which are mainly oceanic, open-water species which lack a corneal membrane covering the eye, and the Myopsida which are mainly near-shore species and which possess a corneal membrane. This membrane has been described as an adaptation preventing ocular damage in shallow, sediment-rich waters of continental shelf areas (Morton and Yonge, 1964). *Ctenopteryx sicula* (Vérany, 1851) (family Ctenopterygidae) is a bathypelagic squid species of cosmopolitan oceanic distribution (Nesis, 1987; Roeleveld *et al.*, 1992). The species lacks an eye-covering corneal membrane and as such is classified in the suborder Oegopsida along with other families of squid exhibiting this characteristic (for example, see Roper *et al.*, 1969; Nesis, 1987). Naef (1923) and Young (1991) have however highlighted a number of morphological features which are at odds with this scheme of classification. *C. sicula* exhibits fusion of nerve axons in the giant fiber system, and has accessory nidamental glands, features which are otherwise found exclusively within the myopsid families Loliginidae and Pickfordioteuthidae. In addition, *C. sicula* has attachments of the fourth buccal connectives similar to those of *Loligo* species, and has suckers on the buccal lappets, exhibits retraction of tentacles into pockets, and has straight, simple funnel locking cartilages, all of which are reminiscent of loliginid morphology (Naef, 1923; Nesis, 1987; Young,

1991). Young (1991) consequently suggested that *C. sicula* could be a squid species related to the loliginids which has become adapted to life in deep water. Here biochemical genetic techniques are used to investigate relationships between *C. sicula* and member species of the families Loliginidae, Histiototeuthidae, Ommastrephidae, and Enoploteuthidae in an attempt to determine to which families it is most closely related, and hence with which suborder the species has closest affinity.

Enzyme electrophoresis is well established as a tool for investigation of taxonomy and systematics (Avice, 1974, 1983; Ferguson, 1980; Ayala, 1983; Richardson *et al.*, 1986; Thorpe and Solé-Cava, 1994), and has been applied to a number of cephalopod problems (Smith *et al.*, 1981; Augustyn and Grant, 1988; Brierley and Thorpe, 1994; Yokowa, 1994; Brierley *et al.*, in press). The theoretical basis for the use of electrophoretic data in taxonomic studies is founded on what has become known as the *molecular clock hypothesis* (Thorpe, 1982; Nei, 1987). This hypothesis holds that between reproductively isolated groups of organisms molecules such as enzymes, the structures of which are under direct genetic control, diverge at a rate that is stochastically related to the evolutionary time since divergence. The majority of cases employing electrophoresis as a taxonomic aid have addressed questions raised at the level of populations, species, or genera. The technique has however also provided meaningful data for comparisons between species from related families (for example, see Solé-Cava *et al.*, 1992, 1994), and is therefore an appropriate means for investigation of the question in hand.

\*Present address: British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, United Kingdom.



## MATERIALS AND METHODS

### Squid samples

*Ctenopteryx sicula* (Vérany, 1851) and the oegopsid squid species *Pyroteuthis margaritifera* (Ruppell, 1844), *Histioteuthis bonellii* (Férussac, 1835), and *Todarodes sagittatus* (Lamarck, 1799) were caught from the northeastern Atlantic Ocean in the region of 21°W, 31°N using a multiple-opening, 8 m<sup>2</sup> rectangular mid-water trawl (RMT8) during RRS *Discovery* cruise 194 in August 1990 (see Herring, 1990). *Sepioteuthis lessoniana* Lesson, 1830, was obtained from Indonesian waters, and *Loligo forbesi* Steenstrup, 1856, and *Loligo vulgaris vulgaris* Lamarck, 1798, were caught around the British Isles. Samples of mantle tissue were dissected from each specimen as soon after capture as possible, and immediately frozen (to -80°C in the case of the *Discovery* samples) prior to transportation to Port Erin Marine Laboratory. Date and location of capture, and species sample sizes, are presented in Table 1.

### Electrophoresis

Mantle tissue samples from *Ctenopteryx sicula* and the other oegopsid species were assayed during September 1990 using a series of standard electrophoretic techniques which have been described in detail elsewhere (Brierley, 1992; Brierley *et al.*, 1993a, b). The majority of the loliginids had been examined previously using the same experimental protocols, but a group of ten were run again concurrently with the *Discovery* samples in order to provide a reference for relative band migration. Gels were scored immediately after optimum stain development, and genotypes assigned accordingly.

### Data analysis

Allele frequency data was analyzed using the FORTRAN program BIOSYS-1 (Release 1.7) (Swofford and Selander, 1981) to calculate unbiased genetic identity (*I*) and distance (*D*) (Nei, 1978) between species pairs, and to construct a dendrogram of *D* using the unweighted pair-

grouping arithmetic mean (UPGMA) cluster analysis algorithm (Sneath and Sokal, 1973). The use of the UPGMA method in conjunction with *D* is generally recognized as the best and most accurate procedure for phylogenetic tree reconstruction from electrophoretic data (Nei, 1987). This is partly because *D* can be subject to large stochastic errors (especially if calculated with small numbers of loci), and the procedure of distance-averaging used in UPGMA reduces this error considerably (Nei, 1987).

## RESULTS

Seventeen enzyme stain systems developed successfully for all species, revealing the presence of 18 common putative enzyme loci. The group of ten loliginids run concurrently with *Ctenopteryx sicula* and the oegopsid species resolved identically here as previously (Brierley *et al.*, 1993a, 1995; Brierley and Thorpe, 1994). A slight reduction in band intensity was apparent, but band migration and definition remained entirely unaffected by frozen (-26°C) storage. Allele frequencies at the 18 clearly resolving enzyme loci are given for each species in Table 2. Inspection of Table 2 reveals *Histioteuthis bonellii*, *Todarodes sagittatus*, and *Pyroteuthis margaritifera* exhibiting very low levels of mean heterozygosity per locus, a phenomenon which is a seemingly recurrent theme in studies of squid biochemical genetics (see review in Brierley *et al.*, 1993a). Although this ubiquitous feature of low intraspecific genetic variability can cause problems for the use of electrophoretic data in squid population studies (see Brierley *et al.*, 1993b, 1995), taxonomic studies benefit because banding patterns remain simple, aiding interpretation and greatly reducing the likelihood of error (Garthwaite *et al.*, 1989). Mean *I* and *D* between all species pairs are given in Table 3. These unbiased estimates are most appropriate for small sample sizes, especially, as here, when observed heterozygosity levels are low. *D* is purported to be stochastically linear with evolutionary time (Nei, 1987; Thorpe, 1989), and varies between a theoretical minimum of zero in comparison of genetically identical individ-

Table 1. Sample size, and date and location of capture of species examined.

Species	Sample size	Date of capture	Location of capture
<i>Ctenopteryx sicula</i>	4	August 1990	Northeastern Atlantic Ocean
<i>Pyroteuthis margaritifera</i>	11	August 1990	Northeastern Atlantic Ocean
<i>Histioteuthis bonellii</i>	14	August 1990	Northeastern Atlantic Ocean
<i>Todarodes sagittatus</i>	2	August 1990	Northeastern Atlantic Ocean
<i>Sepioteuthis lessoniana</i>	5	April 1988	South Alas Strait, Indonesia
<i>Loligo forbesi</i>	20	October 1989	Isle of Man, British Isles
<i>L. vulgaris vulgaris</i>	20	October 1989	Plymouth, United Kingdom



Table 2. Allele frequencies at the 18 clearly resolving enzyme loci.

Locus	Allele	<i>Ctenopteryx sicula</i> N = 4	<i>Loligo forbesi</i> N = 20	<i>L. v. vulgaris</i> N = 20	<i>Sepioteuthis lessoniana</i> N = 5	<i>Pyroteuthis margaritifera</i> N = 11	<i>Histioteuthis bonnellii</i> N = 14	<i>Todarodes sagittatus</i> N = 2
<i>aGpdh</i>	A	0	0	0	0	0	1	0
	B	0.125	0	0	0	0	0	0
	C	0.875	0	0	0	0	0	1
	D	0	0	0.9	0	0	0	0
	E	0	1	0.1	0	0	0	0
	F	0	0	0	1	1	0	0
<i>Mdh-1</i>	A	0	1	1	0	0	0	0
	B	0	0	0	1	0	0	0
	C	0.875	0	0	0	0	0	0
	D	0	0	0	0	1	0	1
	E	0.125	0	0	0	0	1	0
<i>Mdh-2</i>	A	0	0	0	0	0	1	0
	B	1	0	0	1	0	0	0
	C	0	1	1	0	0	0	0
	D	0	0	0	0	0	0	1
	E	0	0	0	0	1	0	0
<i>Me</i>	A	0	0	0	1	0	0	0
	B	0	1	1	0	0	0	0
	C	0	0	0	0	1	0	0
	D	1	0	0	0	0	0	0
	E	0	0	0	0	0	1	0
	F	0	0	0	0	0	0	1
<i>Idh</i>	A	0	0	0	0	0	0	1
	B	0	0	0	0	0	1	0
	C	1	0	0	1	0	0	0
	D	0	0.95	1	0	0	0	0
	E	0	0.05	0	0	1	0	0
<i>Pgdh</i>	A	0	0	0	1	0	0	0
	B	0	0	1	0	0	0	0
	C	0	0.025	0	0	0	0	0
	D	0	0.975	0	0	0	0	0
	E	0	0	0	0	1	0	0
	F	1	0	0	0	0	0	0
	G	0	0	0	0	0	1	0
	H	0	0	0	0	0	0	1
<i>Gopdh</i>	A	0	0	0	0	1	0	0
	B	0	0	0	0	0	0	1
	C	0	1	0	1	0	0	0
	D	0	0	1	0	0	0	0
	E	0	0	0	0	0	1	0
	F	1	0	0	0	0	0	0
<i>Gapdh</i>	A	0	0	0	0	1	0	0
	B	0	0	0	0	0	0	1
	C	0	1	0	0	0	0	0
	D	0	0	0	1	0	0	0
	E	1	0	1	0	0	0	0
	F	0	0	0	0	0	1	0
<i>Sdh</i>	A	0	0	0	0	0	1	0
	B	0	0	0	0	1	0	0
	C	1	0	0	0	0	0	0
	D	0	1	0	0	0	0	0
	E	0	0	0	1	0	0	1
	F	0	0	1	0	0	0	0
<i>Pep-A</i>	A	1	0	0	0	0	0	0
	B	0	1	0	0	0	0	0
	C	0	0	1	0	0	0	0
	D	0	0	0	0	0	0	1
	E	0	0	0	1	0	0	0
	F	0	0	0	0	1	0	0
	G	0	0	0	0	0	1	0

(Continued)



Table 2. Continued.

Locus	Allele	<i>Ctenopteryx sicula</i> N = 4	<i>Loligo forbesi</i> N = 20	<i>L. v. vulgaris</i> N = 20	<i>Sepioteuthis lessoniana</i> N = 5	<i>Pyroteuthis margaritifera</i> N = 11	<i>Histioteuthis bonnellii</i> N = 14	<i>Todarodes sagittatus</i> N = 2
<i>Ck</i>	A	0.125	0	0	0	0	0	0
	B	0.875	0	0	0	0	0	0
	C	0	1	1	0	0	0	0
	D	0	0	0	1	0	0	0
	E	0	0	0	0	1	1	1
<i>Fum</i>	A	0	0	0	0	0	0	1
	B	0	1	0	0.5	0	0	0
	C	1	0	0	0	0	0	0
	D	0	0	0	0	1	0	0
	E	0	0	1	0	0	0	0
<i>Mpi</i>	F	0	0	0	0.5	0	0	0
	G	0	0	0	0	0	1	0
	A	0	0	0	0.5	0	0	0
	B	0	0	0	0	0	0	1
	C	0	0	0	0.5	0	0	0
<i>Gpi</i>	D	0	0	1	0	0	0	0
	E	1	0.975	0	0	0	0	0
	F	0	0.025	0	0	1	1	0
	A	0	0	0	0	0	0	1
	B	0	0	0	0	1	0	0
<i>Ald</i>	C	1	0	0	0	0	0	0
	D	0	1	1	0	0	0	0
	E	0	0	0	1	0	0	0
	F	0	0	0	0	0	1	0
	A	0	1	1	1	0	1	1
<i>Odh</i>	B	1	0	0	0	0	0	0
	C	0	0	0	0	1	0	0
	A	0	1	0	1	0	0	0
	B	1	0	1	0	1	0	0
	C	0	0	0	0	0	1	0
<i>Ldh</i>	D	0	0	0	0	0	0	1
	A	1	0	0	0	0	0	0
	B	0	1	0.825	0	0	0	0
	C	0	0	0	0	1	0	0
	D	0	0	0	1	0	0	0
<i>Sordh</i>	E	0	0	0.175	0	0	0	0
	F	0	0	0	0	0	1	0
	G	0	0	0	0	0	0	1
	A	0	0.125	0	0	0	0	0
	B	0	0.85	0	1	0	0	0
	C	0	0.025	0	0	0	0	0
	D	0	0	1	0	0	0	0
	E	1	0	0	0	0	0	0
	F	0	0	0	0	1	0	0
	G	0	0	0	0	0	0	1
	H	0	0	0	0	0	1	0

Table 3. Matrix of unbiased genetic identity (*I*) (above diagonal) and distance (*D*) (below diagonal) values between all species pairs.

Species	1	2	3	4	5	6	7
1 <i>Ctenopteryx sicula</i>	****	0.056	0.115	0.117	0.057	0.007	0.05
2 <i>Loligo forbesi</i>	2.881	****	0.449	0.253	0.004	0.058	0.056
3 <i>L. vulgaris vulgaris</i>	2.162	0.800	****	0.058	0.056	0.056	0.056
4 <i>Sepioteuthis lessoniana</i>	2.144	1.375	2.845	****	0.057	0.057	0.115
5 <i>Pyroteuthis margaritifera</i>	2.869	5.468	2.877	2.859	****	0.111	0.111
6 <i>Histioteuthis bonnellii</i>	4.949	2.853	2.877	2.859	2.197	****	0.111
7 <i>Todarodes sagittatus</i>	3.003	2.877	2.877	2.165	2.197	2.197	****



uals, and infinity when the pair of taxa under consideration exhibits no common genetic characters. A UPGMA dendrogram of  $D$  (Fig. 1) shows *Ctenopteryx sicula* clustering more closely with the branch encompassing the loliginid species than to the branch containing the oegopsid families Histioeuthidae, Ommastrephidae, and Enoplateuthidae.

## DISCUSSION

In contrast to electrophoretic studies of population structuring, in which large sample sizes are required, taxonomic studies such as the one reported here can be successfully conducted using only very small numbers of individuals from each taxa if large numbers of enzyme loci are studied (see for example Richardson *et al.*, 1986). The comparatively small sample sizes of *Ctenopteryx sicula*, *Todarodes sagittatus*, and *Sepioteuthis lessoniana* are unlikely to be a major source of inaccuracy in calculation of  $D$ , or, consequently, in reconstruction of the phylogenetic tree shown in Fig. 1. Numbers of animals used have negligible effects upon the errors of genetic distance, and even sample sizes as small as one will give acceptable distance estimates (see Nei, 1978; Gorman and Renzi, 1979; Thorpe, 1982). This fact is well illustrated by a reconstructed phylogeny of the family Ommastrephidae (Yokowa, 1994) which is based on electrophoretic data gathered from 16 species with a mean sample size of less than two.

Yeatman and Benzie (1993) used *Sepioteuthis lessoniana* as an outgroup in a study of cryptic speciation in *Loligo* from Australian waters. They reported the species differing completely from all *Loligo* taxa investigated at eight of the 11 loci they resolved, and consequently report-

ed a  $D$  value of  $> 4$  between species. Such a high value is far larger than would normally be expected between members of confamilial genera (Thorpe, 1982). We have assayed six (*Ak*, *Gpi*, *Idh*, *Mdh-1*, *Mdh-2*, and *Mpi*) of these eight loci here and elsewhere (Brierley, 1992; Brierley *et al.*, 1993b, in press) and similarly find alleles unique to *Sepioteuthis lessoniana* at these loci. However this and our previous analyses have included data from a number of other loci in addition to the 11 investigated by Yeatman and Benzi (1993), and our estimate of  $D$  between *Loligo* species and *Sepioteuthis lessoniana* of the order of 1.4 is therefore likely to be the more accurate. This large discrepancy between studies highlights the possibility of substantial interlocus errors on measures of genetic distance (Thorpe, 1979, 1982), and emphasizes the importance of screening large numbers of loci.

The phylogenetic tree reconstructed using UPGMA analysis of  $D$  between all species pairs (Fig. 1) comprises two main branches, one containing member species of the suborder Oegopsida, and the other containing myopsid species. *Ctenopteryx sicula* clusters more closely with the Myopsida, and genetic data presented here are therefore in agreement with morphological evidence (Naef, 1923; Young, 1991) suggesting the species is related to *Loligo*. *C. sicula* clusters with other myopsid species at a level of  $D$  which lies within the practical limits of the measure. The  $D$  value of about 2.9 exhibited here between *C. sicula* and *Loligo forbesi* (see Table 3) is less than the value of  $D = 3$ , beyond which the measure diverges from linearity (Thorpe, 1989). This limit arises because  $D$  appears to suffer a "saturation" effect (Thorpe, 1982): a maximum of one nucleotide substitution per locus can be detected electrophoretically, and practical limitations impose a resolution threshold on




Fig. 1. UPGMA dendrogram of Nei's (1978) unbiased genetic distance ( $D$ ) between all species pairs.



the number of discrete alleles which can be electrophoretically distinguished. Using data gathered from a survey of around 100 loci between species pairs from different major taxa, Thorpe (1982) found an approximate 5% coincidental identity between quite unrelated taxa ( $I$  in the order of 0.05,  $D = -\log_e I$ ). The similarity between *C. sicula* and the loliginid species studied here is clearly greater than this 5% similarity. The level at which the oegospid and myopsid species cluster is high, but this does not detract from the validity of the observation that members of the Myopsida and Oegopsida cluster as two discrete entities, or that *C. sicula* is more closely related to loliginids than to any of the oegospid families studied here. We do not attempt to place an exact time of divergence on any of the branch points in Fig. 1, because the absolute relationship between genetic distance and evolutionary time is questionable (Sarich, 1977; Lessios, 1979; Thorpe, 1982; Smith and Coss, 1984). For most taxonomic purposes however, as here, the problem of absolute nonlinearity of  $D$  with time is unimportant because only relative values are needed (Thorpe, 1982). As long as genetic distance is approximately linear with time, the relative evolutionary relationships among organisms can be determined (Nei, 1987).

*Ctenopteryx sicula* is positioned within the branch of the phylogenetic tree (Fig. 1) containing *Loligo forbesi* and *L. vulgaris vulgaris* beyond the junction at which *Sepioteuthis lessoniana* diverges from the genus *Loligo*. *Sepioteuthis* has been described, on morphological and distributional grounds, as a Tethyan relict genus which split off from all other loliginids before the closing of the Tethys Sea (Brakoniecki, 1986). *C. sicula* would therefore appear to have diverged from the loliginid line at some time before this event. Naef (1923) suggested that *C. sicula* should be considered the earliest independent branch of the stem of the higher Oegopsida. Genetic data from the present study summarized in the phenogram in Fig. 1 go some way to support this assertion, although further electrophoretic analysis of *Pickfordiateuthis pulchella* Voss, 1953, would be necessary before genetic and morphological data could be viewed in complete congruence. It would seem appropriate therefore to adopt the viewpoint of Young (1991), who has proposed that *C. sicula* may be considered as a teuthoid related to loliginids which has become adapted to a deep-water, oceanic life. Young (1991) has also argued that, despite the fact that the name is inappropriate for a squid exhibiting the open-eyed condition, it would be desirable to classify *Ctenopteryx* species with the families Loliginidae and Pickfordiateuthidae in the suborder Myopsida.

The numerous similarities between *Ctenopteryx* and *Bathyteuthis* species have been well described (Roper, 1969; Nesis, 1987), and more recent analyses of comparative beak morphology have further highlighted the affinity

among *Loligo*, *Ctenopteryx*, and *Bathyteuthis* species (Roper and Clarke, unpub. data). It would be of additional interest to examine *Bathyteuthis* species electrophoretically, as it remains possible that these species similarly fail to comply with the broad categorization of an open-eyed, open-ocean suborder, contrasting entirely with a distantly related shelf-inhabiting group. Our data suggest that the character of the presence or absence of a corneal membrane may not be an infallible guide for distinguishing the two major systematic groupings of squid.

## ACKNOWLEDGMENTS

We should like to thank Dr. Peter Herring for providing the opportunity to collect samples during RRS *Discovery* cruise 194, and for invaluable help with squid identification at sea. We also thank the scientists, Captain and crew of RRS *Discovery* cruise 194, the Captain and crew of Liverpool University's RV *Cuma*, Dr. A. Ghofar and Mr. P. Pascoe for assistance with sample collection, and Professor T. Norton for provision of laboratory facilities at Port Erin Marine Laboratory. This study was supported by grant number GR3/6849 awarded by the Natural Environment Research Council to Dr. J. P. Thorpe.

## LITERATURE CITED

- Augustyn, C. J. and W. S. Grant. 1988. Biochemical and morphological systematics of *Loligo vulgaris vulgaris* Lamarck and *Loligo vulgaris reynaudi* D'Orbigny nov. comb. (Cephalopoda: Myopsida). *Malacologia* 29:215-233.
- Awise, J. C. 1974. Systematic value of electrophoretic data. *Systematic Zoology* 23:465-481.
- Awise, J. C. 1983. Protein variation and phylogenetic reconstruction. In: *Protein Polymorphism: Adaptive and Taxonomic Significance*, G. S. Oxford and D. Rollinson, eds. pp. 103-130. Academic Press, London.
- Ayala, F. J. 1983. Enzymes as taxonomic characters. In: *Protein Polymorphism: Adaptive and Taxonomic Significance*, G. S. Oxford and D. Rollinson, eds. pp. 3-26. Academic Press, London.
- Brakoniecki, T. F. 1986. *A Generic Revision of the Family Loliginidae (Cephalopoda: Myopsida) Based Primarily on the Comparative Morphology of the Hectocotylus*. Doctoral dissertation, University of Miami, Miami, Florida. 163 pp.
- Brierley, A. S. 1992. *Aspects of Genetic Diversity and Population Structure in Squid*. Doctoral dissertation, University of Liverpool. 154 pp.
- Brierley, A. S. and J. P. Thorpe. 1994. Biochemical genetic evidence supporting the taxonomic separation of *Loligo gahi* from the genus *Loligo*. *Antarctic Science* 6(2):143-148.
- Brierley, A. S., P. G. Rodhouse, J. P. Thorpe, and M. R. Clarke. 1993a. Genetic evidence of population heterogeneity and cryptic speciation in the ommastrephid squid *Martialia hyadesi* from the Patagonian Shelf and Antarctic polar frontal zone. *Marine Biology* 116(4):593-602.
- Brierley, A. S., J. P. Thorpe, M. R. Clarke, and H. R. Martins. 1993b. A preliminary biochemical genetic investigation of the population structure of *Loligo forbesi* Steenstrup, 1856 from the British Isles and the Azores. In: *Recent Advances in Cephalopod Fisheries Biology*, T. Okutani, R. K. O'Dor, and T. Kubodera, eds. pp. 61-



69. University Press, Tokyo.
- Brierley, A. S., J. P. Thorpe, G. J. Pierce, M. R. Clarke, and P. R. Boyle. 1995. Genetic variation in the neritic squid *Loligo forbesi* (Myopsida: Loliginidae) in the northeast Atlantic. *Marine Biology* 122:79-86.
- Brierley, A. S., A. L. Allcock, J. P. Thorpe, and M. R. Clarke. In press. Biochemical genetic evidence supporting the taxonomic separation of *Loligo edulis* and *Loligo chinensis* (Cephalopoda: Teuthoidea) from the genus *Loligo*. *Marine Biology*.
- Ferguson, A. 1980. *Biochemical Systematics and Evolution*. Blackie, Glasgow. 194 pp.
- Garthwaite, R. L., C. J. Berg, Jr., and J. Harrigan. 1989. Population genetics of the common squid *Loligo pealei* Le Seur, 1821, from Cape Cod to Cape Hatteras. *Biological Bulletin* 177:287-294.
- Gorman, G. C. and J. Renzi. 1979. Genetic distance and heterozygosity estimates in electrophoretic studies: effects of sample size. *Copeia* 1979:361-364.
- Herring, P. J. 1990. *RRS Discovery Cruise 194 04 Aug-02 Sep 1990. Comparative Benthic Biology at 31°N 20°W (Madeira Abyssal Plain) and Associated Midwater Studies*. Institute of Oceanographic Sciences Deacon Laboratory, Cruise Report 219. 57 pp.
- Lessios, H. A. 1979. Use of Panamanian sea urchins to test the molecular clock. *Nature* 280:599-601.
- Morton, J. E. and C. M. Yonge. 1964. Classification and structure of the Mollusca. In: *Physiology of Mollusca*, K. M. Wilbur and C. M. Yonge, eds. pp. 1-58. Academic Press, New York.
- Naef, A. 1921-1923. *Fauna and Flora of the Bay of Naples*. 35. *Cephalopoda*. English translation by the Israel Program for Scientific Translation in 1972, Jerusalem. 917 pp.
- Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583-590.
- Nei, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York. 512 pp.
- Nesis, K. N. 1987. *Cephalopods of the World*. T. F. H. Publications, Neptune City, New Jersey. 351 pp.
- Richardson, B. J., P. R. Baverstock, and M. Adams. 1986. *Allozyme Electrophoresis - a Handbook for Animal Systematics and Population Studies*. Academic Press, Sydney. 410 pp.
- Roeleveld, M. A. C., M. R. Lipinski, C. J. Augustyn, and B. A. Stewart. 1992. The distribution and abundance of cephalopods on the continental-slope of the eastern south-Atlantic. *South African Journal of Marine Science* 12:739-752.
- Roper, C. F. E. 1969. Systematics and zoogeography of the worldwide bathypelagic squid *Bathyteuthis* (Cephalopoda, Oegopsida). *Bulletin of the United States National Museum* 291, 210 pp.
- Roper, C. F. E., R. E. Young, and G. L. Voss. 1969. An illustrated key to the families of the order Teuthoidea (Cephalopoda). *Smithsonian Contributions to Zoology* 13, 32 pp.
- Sarich, V. M. 1977. Rates, sample sizes and the neutrality hypothesis for electrophoresis in evolutionary studies. *Nature* 265:24-48.
- Smith, D. G. and R. G. Coss. 1984. Calibrating the molecular clock: estimates of ground squirrel divergence made using fossil and geological time markers. *Molecular Biology and Evolution* 1:249-259.
- Smith, P. J., P. E. Roberts, and R. J. Hurst. 1981. Evidence for two species of arrow squid in the New Zealand fishery. *New Zealand Journal of Marine and Freshwater Research* 15:247-253.
- Sneath, P. H. A. and R. R. Sokal. 1973. *Numerical Taxonomy. The Principles and Practise of Numerical Classification*. Freeman, San Francisco, California. 573 pp.
- Swofford, D. L. and R. B. Selander. 1981. BIOSYS-1: A FORTRAN program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *Journal of Heredity* 72:281-283.
- Solé-Cava, A. M., N. Boury-Esnault, J. Vacelet, and J. P. Thorpe. 1992. Biochemical genetic divergence and systematics in sponges of the genera *Corticium* and *Oscarella* (Demospongiae: Homoscleromorpha) in the Mediterranean Sea. *Marine Biology* 113:299-304.
- Solé-Cava, A. M., C. A. M. Russo, M. E. Araujo, and J. P. Thorpe. 1994. Cladistic and phenetic analysis of allozyme data for nine species of sea anemones of the family Actiniidae (Cnidaria: Anthozoa). *Biological Journal of the Linnean Society* 52(3):225-239.
- Thorpe, J. P. 1979. Enzyme variation and taxonomy; the estimation of sampling errors in measurement of interspecific genetic similarity. *Biological Journal of the Linnean Society* 11:369-386.
- Thorpe, J. P. 1982. The molecular clock hypothesis: biochemical evolution, genetic differentiation and systematics. *Annual Review of Ecology and Systematics* 13:139-168.
- Thorpe, J. P. 1989. Possible effects of interprotein variation in mean rate of amino acid substitution on the relationship of genetic distance with time since evolutionary divergence. *Biological Journal of the Linnean Society* 37:335-344.
- Thorpe, J. P. and A. M. Solé-Cava. 1994. The use of electrophoresis in invertebrate systematics. *Zoological Scripta* 23(1):3-18.
- Yeatman, J. and J. A. H. Benzie. 1993. Cryptic speciation in *Loligo* from northern Australia. In: *Recent Advances in Cephalopod Fisheries Biology*, T. Okutani, R. K. O'Dor, and T. Kubodera, eds., pp. 641-652. Tokai University Press, Tokyo.
- Yokawa, K. 1994. Allozyme differentiation of sixteen species of ommastrephid squid (Mollusca, Cephalopoda). *Antarctic Science* 6(2):210-204.
- Young, J. Z. 1991. *Ctenopteryx* the comb-fin squid is related to *Loligo*. *Bulletin of Marine Science* 49(1-2):148-161.

Date of manuscript acceptance: 5 February 1996